



4TH INTERNATIONAL PROPOLIS CONFERENCE

PROPOLIS - IMMUNITY - COMMUNITY - SUSTAINABILITY

13-15th February 2025
Ribeirão Preto - Brazil

BOOK OF
ABSTRACTS

Dados Internacionais de Catalogação na Publicação (CIP)
Instituto Brasileiro de Informação em Ciência e Tecnologia, Brasil

4th International Conference Propolis (02.: 2025: Ribeirão Preto, SP) Anais / 4th International Conference Propolis, 13 a 15 de fevereiro de 2025 em Ribeirão Preto, São Paulo; editores: Yara Sbrolin Roldão-Sbordoni ... [et al.]. – Ribeirão Preto: USP, 2025. 101 p.

ISBN: 978 65 01 39259 2

1. Abelha - Congressos. I. Roldão-Sbordoni, Yara Sbrolin. II. Almeida-Dias, Joyce Mayra Volpini. III. Berretta, Andresa. IV. Universidade de São Paulo. V. Título.

CDD-595.79906

CDU: 595.799



Abstracts of papers presented at the



Campus USP Ribeirão Preto, Brazil, 13 – 15th February 2025.

Staff Commission:

Adriel Santos

Aline Takeuchi

Bianca Mariano

Cristiana Cunha

Herena Oliveira

Jéssica Aparecida de Lima

Joyce Mayra V. de Almeida-Dias

Michelangelo Rodrigues

Rafaela Depintor

Thaís Faggion

Thiago Depintor

Yara S. Roldão-Sbordoni

Scientific Commission:

Andressa A. Berretta

Antônio Salatino

Badiaa Lyoussi

David de Jong

James Fearnley

Marcelo Augusto Duarte Silveira

Miguel Vilas-Boas

Trong Tran

Vassya Bankova

Edited by:

Almeida-Dias, J.M.V. & Roldão-Sbordoni, Y.S.



ORGANIC PROPOLIS FROM URUGUAY: DEVELOPING A METHODOLOGY TO ANALYZE ITS VOLATILE COMPONENTS

Manuel Minteguiaga^{1,2}, Francisco Santurión³, Eduardo Dellacassa²

¹*Espacio de Ciencia y Tecnología Química, CENUR Noreste, Universidad de la República, 45000, Tacuarembó, Uruguay.*

²*Laboratorio de Biotecnología de Aromas, Departamento de Química Orgánica, Facultad de Química, Universidad de la República, 11800, Montevideo, Uruguay.*

³*Independent organic beekeeper.*

E-mails: manuel.minteguiaga@pedeciba.edu.uy, edellac@fq.edu.uy

Propolis volatile components (PVCs) not only contribute to the product's pleasant aroma but also exhibit a wide range of biological activities, including antimicrobial, immunomodulatory, and cytotoxic effects against human cancer cells [1]. Additionally, the volatile composition of propolis serves as an indicator of its botanical and geographical origin, making it a key parameter for quality control [1]. In this ongoing project, we outline the initial steps in developing an analytical methodology to study PVCs. To date, only one previous study has reported PVCs from Uruguayan samples, using static headspace sampling, but without specifying the origin or production conditions [2]. In the present work, we analyze samples from an organic apiary in Rocha Province, eastern Uruguay, using sampling best practices in accordance with official recommendations [3]. Headspace sampling of 0.03 g of ground propolis was conducted at 50°C for 30 minutes [2] using two different SPME fibers (1, DVB/PDMS/Carboxen; 2, Polyacrylate). The extracted PVCs were then directly desorbed in the injector port of a GC-MS instrument. Two stationary phases (Rxi-5MS and Stabilwax-MS) were selected for conventional analysis, using an optimized oven program. Identification of PVCs was carried out by comparing mass spectra with commercial libraries and calculating linear retention indices (LRIs) using a C₈-C₂₀ alkane solution [4]. Additionally, separation of chiral monoterpenes was achieved via eGC-MS with an Rt- β DEXsm column (stationary phase composed of a modified β -cyclodextrin as chiral selector). Over 100 PVCs were detected, predominantly phenylpropanoids, benzenoids and terpenic compounds. Some key PVCs identified include α -pinene, benzyl alcohol, terpinen-4-ol, guaiacol, benzyl benzoate, spathulenol and β -selinene. The detection of *trans*-nerolidol suggests *Baccharis dracunculifolia* (Asteraceae) as the botanical origin of the samples, like Brazilian Green Propolis [5]. Further studies on the chiral patterns of selected PVCs could enhance quality control, define potential markers and support origin certification of this product in Uruguay.

Key Words: Propolis, Volatile Components, Organic Production, HS-SPME, GC-MS, eGC-MS.

Acknowledgements: PEDECIBA-Program for the Development of Basic Sciences and SNI-National System of Researchers (Uruguay) are acknowledged for financial support for conducting this research.